arXiv:1701.02602v3 [math.NT] 21 Feb 2017

IS THE QUARTIC DIOPHANTINE EQUATION $A^4 + hB^4 = C^4 + hD^4$ SOLVABLE FOR ANY INTEGER h?

FARZALI IZADI AND MEHDI BAGHALAGHDAM

Abstract. The Diophantine equation $A^4 + hB^4 = C^4 + hD^4$, where h is a fixed arbitrary positive integer, has been investigated by some authors. Currently, by computer search, the integer solutions of this equation are known for all positive integer values of $h \leq 5000$ and $A, B, C, D \leq 100000$, except for some numbers, while a solution of this Diophantine equation is not known for arbitrary positive integer values of h. Gerardin and Piezas found solutions of this equation when h is given by polynomials of degrees 5 and 2 respectively. Also Choudhry presented some new solutions of this equation when h is given by polynomials of degrees 2, 3, and 4.

In this paper, by using the elliptic curves theory, we study this Diophantine equation, where h is a fixed arbitrary rational number. We work out some solutions of the Diophantine equation for certain values of h, in particular for the values which has not already been found a solution in the range where $A, B, C, D \leq 100000$ by computer search. Also we present some new parametric solutions for the Diophantine equation when h is given by polynomials of degrees 3, 4. Finally We present two conjectures such that if one of them is correct, then we may solve the above Diophantine equation for arbitrary rational number h.

1. Introduction

The Diophantine equation (DE)

$$A^4 + hB^4 = C^4 + hD^4, \tag{1.1}$$

where h is a fixed arbitrary positive integer, has been investigated by some authors.

The numerical solutions for 75 integer values of $h \leq 101$ was given by Choudhry [A.C]. Then these solutions were first extended by Piezas [T.P] for all positive integer values of $h \leq 101$, and finally by Tomita [S.T] for all positive integer values of $h \leq 1000$ except $h = 967$. The lost solution for $h = 967$ was supplied by Bremner. Currently, by computer search, the small solutions of this DE are known for all positive integers $h \leq 5000$, and $A, B, C, D \leq 100000$ except for

2010 *Mathematics Subject Classification.* 11D45, 11D72, 11D25, 11G05 and 14H52.
Key words and phrases. Quaratic Diophantine equation, Biquadratics, Elliptic curves.

$h = 1198, 1787, 1987,$

$2459, 2572, 2711, 2797, 2971,$

$3086, 3193, 3307, 3319, 3334, 3347, 3571, 3622, 3623, 3628, 3644,$
$3646, 3742, 3814, 3818, 3851, 3868, 3907, 3943, 3980,$

$4003, 4006, 4007, 4051, 4054, 4099, 4231, 4252, 4358, 4406, 4414,$
$4418, 4478, 4519, 4574, 4583, 4630, 4643, 4684, 4870, 4955, 4999.$

We will work out some of these cases.

Gerardin and Piezas found solutions of this equation when h is given by polynomials of degrees 5, and 2, respectively, see [T.P] and [S.T]. Also Choudhry presented several new solutions of this equation when h is given by polynomials of degrees 2, 3, and 4, see [A.C2].

In this paper, we used elliptic curves theory to study the DE (1.1).

2. The first method for solving the DE $A^4 + hB^4 = C^4 + hD^4$

Our main result in this section is the following:

Main Theorem 2.1. *Consider the DE* (1.1)*, where h is a fixed arbitrary rational number.*
Then there exists a cubic elliptic curve of the form
$E(h) : Y^2 = X^3 + FX^2 + GX + H$, where the coefficients F, G, and H, are all functions of h. If the elliptic curve $E(h)$ or its counterpart $E(h)_t$ resulting from $E(h)$ by switching h to ht^4 has positive rank, depending on the value of h and an appropriate rational number t, then the DE (1.1) *has infinitely many integers solutions. By taking $h = \frac{v}{u}$, this also solves DE of the form $uA^4 + vB^4 = uC^4 + vD^4$ for appropriate integer values of u and v.*

Proof 2.1. *Let: $A = m - q$, $B = m + p$, $C = m + q$, and $D = m - p$, where all variables are rational numbers. By substituting these variables in the DE* (1.1) *we get*

$$-8m^3q - 8mq^3 + 8hm^3p + 8hmp^3 = 0. \tag{2.1}$$

Then after some simplifications and clearing the case $m = 0$ we obtain

$$m^2(hp - q) = -hp^3 + q^3. \tag{2.2}$$

We may assume that $hp-q=1$ *and* $m^2=-hp^3+q^3$.
By plugging $q=hp-1$ *into the equation* (2.2) *and some simplifications we obtain the equation*

$$m^2=(h^3-h)p^3-(3h^2)p^2+(3h)p-1. \tag{2.3}$$

By multiplying both sides of this equation in $(h^3-h)^2$ *and letting*

$$X=(h^3-h)p \qquad Y=(h^3-h)m, \tag{2.4}$$

we get the elliptic curve

$$Y^2=X^3-(3h^2)X^2+(3h(h^3-h))X-(h^3-h)^2. \tag{2.5}$$

By Letting $X=Z+h^2$ *in* (2.5), *we get the simple elliptic curve*

$$E(h):Y^2=Z^3-(3h^2)Z-(h^4+h^2). \tag{2.6}$$

If for a given h, *the above elliptic curve* $E(h)$ *or its counterpart* $E(h)_t$ *resulting from* $E(h)$ *by switching* h *to* ht^4 *has positive rank, then by calculating* m, p, q, A, B, C, D, *from the relations* (2.4), $q=hp-1$, $A=m-q$, $B=m+p$, $C=m+q$, $D=m-p$, *after some simplifications and canceling the denominators of* A, B, C, D, *we obtain infinitely many integer solutions for the DE* (1.1). *Now the proof of the main theorem is completed.*

Although, we were able to find an appropriate t such that $E(h)_t$ has positive rank in the case of rank zero $E(h)$ for many values of h, the proof for arbitrary h seems to be difficult at this point. For this reason, we state it as a conjecture.

Conjecture 2.2. *Let* h *be an arbitrary fixed rational number. Then there exists at least a rational number* t *such that the rank of the elliptic curve*

$$E(h)_t:Y^2=Z^3-(3h^2t^8)Z-(h^4t^{16}+h^2t^8), \tag{2.7}$$

is positive.

Remark 1. If h is a large number, for example $h=7000$, we may divide h by 10^4 to get an elliptic curve of positive rank with small coefficients, then

solve the DE (1.1) for $h' = \frac{7}{10}$ and finally get a solution for the main case of $h = 7000$, by multiplying both sides of the DE (1.1) by an appropriate number. As another example, if $h = 9317 = 7.11^3$, we may first solve the DE (1.1) for $h' = \frac{7}{11}$.

Remark 2. Note that by substituting the relation $p = \frac{q+1}{h}$, in the equation $m^2 = -hp^3 + q^3$, multiplying both side of the equation by $(\frac{h^2-1}{h^2})^2$, and letting

$$X' = (\frac{h^2-1}{h^2})q \qquad Y' = (\frac{h^2-1}{h^2})m, \tag{2.8}$$

we get another elliptic curve

$$Y'^2 = X'^3 - \frac{3}{h^2}X'^2 - \frac{3(h^2-1)}{h^4}X' - (\frac{h^2-1}{h^3})^2. \tag{2.9}$$

Now, if we set $Y' = \frac{Y}{h^3}$, $X' = \frac{X+1-h^2}{h^2}$, then the elliptic curve (2.9), transforms to the elliptic curve (2.5). This means that two elliptic curves (2.9) and (2.5) are isomorphic.

3. Application to examples

3.1. **Example:** $A^4 + B^4 = C^4 + D^4$**.**
i.e., sums of two biquadrates in two different ways.

$h = 16$, here $h = 1$ replaced by $h = 2^4$.

$E(16)$: $Y^2 = X^3 - 768X^2 + 195840X - 16646400$.

Rank=1.

Generator: $P = (X, Y) = (340, 680)$.

Points: $2P = (313, -275)$, $3P = (\frac{995860}{729}, \frac{-727724440}{19683})$, $4P = (\frac{123577441}{302500}, \frac{305200800239}{166375000})$.

$(p', m', q') = (\frac{313}{4080}, \frac{-55}{816}, \frac{58}{255})$,
$(p'', m'', q'') = (\frac{2929}{8748}, \frac{-1070183}{118098}, \frac{9529}{2187})$,
$(p''', m''', q''') = (\frac{123577441}{1234200000}, \frac{305200800239}{678810000000}, \frac{46439941}{77137500})$.

Solutions:

$$1203^4 + 76^4 = 653^4 + 1176^4,$$

$$1584749^4 + 2061283^4 = 555617^4 + 2219449^4,$$

$$103470680561^4 + 746336785578^4 = 713872281039^4 + 4744664153 78^4.$$

3.2. **Example:** $A^4 + 206B^4 = C^4 + 206D^4$**.**

$h = \frac{103}{8}$.

$E(\frac{103}{8})$: $Y^2 = X^3 - \frac{31827}{64}X^2 + \frac{335615715}{4096}X - \frac{1179689238225}{262144}$.

Rank=1.

Generator: $P = (X, Y) = (\frac{2131205}{32}, \frac{8767168835}{512})$.

$(p, m, q) = (\frac{6819856}{217227}, \frac{1753433767}{217227}, \frac{850373}{2109})$.

Solution:

$$3331690696^4 + 206.(1760253623)^4 = 3682044372^4 + 206.(1746613911)^4.$$

Remark 3. By searching, Noam Elkies found the smallest solution to this DE as $A, B, C, D = 3923, 1084, 4747, 506$.

Remark 4. No computer research has come up with a solution for the following equations in the range of $A, B, C, D \leq 100000$, see [S.T].

3.3. **Example:** $A^4 + 2572B^4 = C^4 + 2572D^4$**.** $h = 2572$.

$E(2572)$: $Y^2 = X^3 - (3.2572^2)X^2 + (3.2572.(2572^3 - 2572))X - (2572^3 - 2572)^2$.

Rank=2.
Generators: $P_1 = (X, Y) = (\frac{60035809}{9}, \frac{302757191}{27})$,

$P_2 = (X', Y') = (\frac{3435573760731933430513}{381659437643236}, \frac{27488556048550361767336062809879}{7456139229698648679016})$.

$(p, m, q) = (\frac{23333}{59513508}, \frac{117667}{178540524}, \frac{194}{23139})$,

$(p', m', q') = (\frac{1558040235953533}{2944884220855208976}, \frac{12466120460409195830562539}{57531570296354773207287456}, \frac{413061923023825}{1144978312929708})$.

Solutions:

$1379237^4 + 2572.(187666)^4 = 1614571^4 + 2572.(47668)^4$,

$828894607040254305529486 1^4 + 2572.(12496558499611049062325037)^4 =$

$33221186991220934716419939^4 + 2572.(12435682421207342598800041)^4$.

3.4. **Example:** $A^4 + 967B^4 = C^4 + 967D^4$**.**
$h = 967$.

$E(967)$: $Y^2 = X^3 - (3.967^2)X^2 + (3.967.(967^3 - 967))X - (967^3 - 967)^2$.

Rank=1.

Generator: $P = (X, Y) = (\frac{238501273696}{245025}, \frac{900632541139856}{121287375})$.

$(p, m, q) = (\frac{2129475658}{1978205172075}, \frac{8041361974463}{979211560177125}, \frac{83761933}{2045713725})$.

Solution:

$320525436849 82^4 + 967.(9095452425173)^4 =$
$481352676339 08^4 + 967.(6987271523753)^4$.

3.5. **Example:** $A^4 + 2797B^4 = C^4 + 2797D^4$**.** $h = 2797$.

$E(2797)$: $Y^2 = X^3 - (3.2797^2)X^2 + (3.2797.(2797^3 - 2797))X - (2797^3 - 2797)^2$.

Rank=1.

Generator: $P=(X,Y)=(\frac{18256234369}{2304},\frac{3411597220289}{110592})$.

$(p,m,q)=(\frac{6527077}{18024671232},\frac{1219734437}{865184219136},\frac{231563137}{18024671232})$.

Solution:

$$9895296139^4+2797.(1533034133)^4=12334765013^4+2797.(906434741)^4.$$

3.6. **Example:** $A^4+4999B^4=C^4+4999D^4$**.** $h=4999$.

$E(4999)$: $Y^2=X^3-(3.4999^2)X^2+(3.4999.(4999^3-4999))X-(4999^3-4999)^2$.

Rank=1.

Generator:$P=(X,Y)=(\frac{3893205338601790029309458312 5}{1502165941669975655844},\frac{5196399152934711936473537677081074562 0625}{5822062544578471664296206412 4328})$.

$p=\frac{62291285417628640468951333}{3002529524556498008097092135 04}$,

$m=\frac{83142386446955390983576602833297192993}{1163713954563345629515072366456370662924 8}$,

$q=\frac{2228682405896333845684837}{6006260301173230662326649 6}$.

Solution:

$A=348665208625932834629908938859838853613,$
$B=855566587295531794454218137167252471 39,$
$C=514949981519843616597062144526433239599,$
$D=807281141643576025217313919498691388 47.$

3.7. **Example:** $A^4+2459B^4=C^4+2459D^4$**.** $h=2459$.

$E(2459)$: $Y^2=X^3-(3.2459^2)X^2+(3.2459.(2459^3-2459))X-(2459^3-2459)^2$.

Rank=1.

Generator: $P = (X, Y)$, where

$X = \frac{24559401683341754492990688766624698 64}{40376478184303184669307544 1721}$,

$Y = \frac{77533931979870341623288869395598504407034176570 0696}{25656218923273051801944840767617022765585 2269}$.

$p = \frac{24979049718614477718664248135 2977}{61060742310948941627623759438304 2485}$,

$m = \frac{78858759133309948762498850076890260788277234103}{38799515034381987144304153520255709273053054862 4665}$,

$q = \frac{14751563526801914704010846945 62}{24831534083346458571624139665 8415}$.

Solution:

$A = 22260874794587190305086350086900354360367782159 59,$
$B = 23758185656414032713676183058169872700517662575 6,$
$C = 23838049977253389280336327088438159576133326841 65,$
$D = 79864338297520429611764130427918205428622157550.$

4. The second method for solving the DE $A^4 + hB^4 = C^4 + hD^4$

In this section, we wish to look at the equation from a different perspective. To this end we take $X = Z^2 + h^2$ (2.5) to get the six degree curve

$$Y^2 = Z^6 - 3h^2Z^2 - (h^4 + h^2). \tag{4.1}$$

This curve can be considered as an quatic elliptic curve of (h, Y) letting Z as a parameter, i.e.,

$$Y^2 = -h^4 - (3Z^2 + 1)h^2 + Z^6. \tag{4.2}$$

Next we use theorem 2.17. of [L.W] to transform this quartic to a cubic elliptic curve of the form

$$E'(Z) : Y'^2 = X'^3 - (3Z^2 + 1)X'^2 + (4Z^6)X' - (12Z^8 + 4Z^6). \tag{4.3}$$

with the inverse transformation

$$h = \frac{2Z^3(X' - (3Z^2 + 1))}{Y'} \qquad Y = -Z^3 + \frac{h^2X'}{2Z^3}. \tag{4.4}$$

Now by taking an appropriate rational value for Z such that the rank of the elliptic curve (4.3) is positive, we obtain an infinitely many rational points on (4.3) and consequently an infinite set of rational values for h (also for Y)), which is denoted by $H(Z)$. Then for every h obtained in this way, we can find a solution for the (2.5) and finally a solution for the main DE by using all the necessary transformations namely
$X = Z^2 + h^2$, $m = \frac{Y}{h^3-h}$, $p = \frac{X}{h^3-h}$, $h = \frac{q+1}{p}$, $A = m - q$, $B = m + p$, $C = m + q$, $D = m - p$.

To get infinitely many solutions one can use the Richmond method [H.R]. The following examples clarify this idea better
$Z_1 = 3$.
$E'(3)$: $Y'^2 = X'^3 - 28X'^2 + 2916X' - 81648$.
Rank=1.
Generator: $P = (X', Y') = (108, 1080)$.
$H(Z_1) = \{4, \frac{2^3.3^3}{197}, \frac{-2^2.251.395449}{11.13.61.653}, \cdots\}$.

$Z_2 = 4$.
$E'(4)$: $Y'^2 = X'^3 - 49X'^2 + 16384X' - 802816$.
Rank=1.
Generator: $P = (X', Y') = (202, 2958)$.
$H(Z_2) = \{\frac{2^6.3}{29}, \frac{2^9.3.29}{61121}, \frac{-2^6.3^2.19.3571.18131}{5.29.97.7746413}, \cdots\}$.

$Z_3 = 6$.
$E'(6)$: $Y'^2 = X'^3 - 109X'^2 + 186624X' - 20342016$.
Rank=1.
Generator: $P = (X', Y') = (\frac{621}{4}, \frac{-24975}{8})$.
$H(Z_3) = \{\frac{-2^2.5.317}{3^3.37}, \frac{2^3.5^4.408841}{17.37^2.12757}, \frac{-2^2.5.10193.249587558933}{3^3.7.37.5101.181680953}, \cdots\}$.

Having seen the examples, the natural question arises:
Does the set of natural numbers $\mathbb{N}$ contained in $\bigcup_{t\in\mathbb{Q}^*} t^4(\bigcup_{Z\in\Omega} H(Z))$?,
where
$\Omega = \{Z \in \mathbb{Q} \mid E'(Z)\ has\ positive\ rank\}$.
We state this as the second conjecture:

Conjecture 4.1. *With the above notations one has* $\mathbb{N} \subset \bigcup_{t\in\mathbb{Q}^*} t^4(\bigcup_{Z\in\Omega} H(Z))$.

5. Application to examples

Now we are going to work out some examples.

5.1. **Example:** $h = 108$**.**
$Z = \frac{5}{3}$.
$E'(\frac{5}{3})$: $Y'^2 = X'^3 - \frac{-28}{3}X'^2 + \frac{62500}{729}X' - \frac{1750000}{2187}$.
Rank=1.
Generator: $P = (X', Y') = (\frac{2500}{81}, \frac{109000}{729})$.
$(m, p, q) = (\frac{5}{4}, \frac{123}{28}, \frac{34}{7})$.
$h = \frac{4}{3}$.
Solution:

$$303^4 + 108(158)^4 = 513^4 + 108(88)^4.$$

Note: $108 = 4.27$.

5.2. **Example:** $h = 492$**.**
$Z = \frac{4}{3}$.
$E'(\frac{4}{3})$: $Y'^2 = X'^3 - \frac{19}{3}X'^2 + \frac{16384}{729}X' - \frac{311296}{2187}$.
Rank=1.
Generator: $P = (X', Y') = (\frac{586}{81}, \frac{5986}{729})$.
$(m, p, q) = (\frac{56908}{11033}, \frac{-238251}{44132}, \frac{-42025}{11033})$.
$h = \frac{64}{123}$.
Solution:

$$42476^4 + 492(395732)^4 = 1863532^4 + 492(59532)^4.$$

Note: $492 = 123.4$.

5.3. **Example** $h = 12256974$**.**
$Z = \frac{3}{2}$.
$E'(\frac{3}{2})$: $Y'^2 = X'^3 - \frac{31}{4}X'^2 + \frac{729}{16}X' - \frac{22599}{64}$.
Rank=1.
Generator: $P = (X', Y') = (\frac{135}{4}, \frac{351}{2})$.
$2P = (\frac{665}{64}, \frac{10309}{512})$.
$h = \frac{54}{61}$.
$(m, p, q) = (\frac{145851}{12880}, \frac{-306037}{19320}, \frac{-48373}{3220})$.
Solutions:

$$62099769^4 + 12256974(174521)^4 = 8718303^4 + 12256974(1049627)^4.$$

Note: $54.61^3 = 12256974$.
For the point 3P, we obtain $h = \frac{-805}{3977}$, then as two more examples, we obtain solutions for the two cases $h' = 3977.805^3$, $h'' = 3977^3.805$.

6. Parametric solutions of $A^4 + hB^4 = C^4 + hD^4$

Let: $A = m-q$, $B = m+p$, $C = m+q$, and, $D = m-p$, where all variables are rational numbers. By substituting these variables in the DE (1.1), and some simplification, we get

$$p(hm^2 + hp^2) = q(m^2 + q^2). \tag{6.1}$$

We may assume that

$$p = m^2 + q^2, \tag{6.2}$$

and

$$q = h(m^2 + p^2). \tag{6.3}$$

By substituting $p = m^2 + q^2$, in the relation $h = \frac{q}{m^2+p^2}$, we get:

$$h = \frac{q}{m^2 + (m^2 + q^2)^2} = \frac{q}{m^2 + m^4 + q^4 + 2m^2q^2}. \tag{6.4}$$

Then using the reciprocal of h, we conclude that

$$h = \frac{m^2 + (m^2 + q^2)^2}{q} = \frac{m^2 + m^4 + q^4 + 2m^2q^2}{q}, \tag{6.5}$$

the DE (1.1) has a parametric solution:
$A = m + m^2 + q^2$,
$B = m - q$,
$C = m - m^2 - q^2$,
$D = m + q$.

Example 6.1. $m = kq$;
$h = k^2q + (k^2 + 1)^2q^3$;
$A = k + k^2q + q$;
$B = k - 1$;
$C = k - k^2q - q$;
$D = k + 1$;

Remark 5. The above example provides parametric solutions for the DE (1.1) when h is given by polynomials of the degrees 3, and 4 as follows.

$m = q$;
$h = q + 4q^3$;
$A = 1 + 2q$;
$B = 0$;
$C = 1 - 2q$;
$D = 2$;

If we let $2q = p$, this recovers the third and the second parametric solutions of the table 1 in [A.C2]. Of course by replacing p with $2p$ in the second parametric solution of the table 1 in [A.C2], and dividing both sides of this equation by 16, we get the third parametric solution of the table 1.!

$h = 8p(p^2 + 1)$;
$A = p + 1$;
$B = 0$;
$C = p - 1$;
$D = 1$;

$m = 2q$;
$h = 4q + 25q^3$;
$A = 2 + 5q$;
$B = 1$;
$C = 2 - 5q$;
$D = 3$;

$m = 3q$;
$h = 9q + 100q^3$;
$A = 3 + 10q$;
$B = 2$;
$C = 3 - 10q$;
$D = 4$;

$q = 2$, $m = 2k$;
$h = 8k^4 + 18k^2 + 8$;
$A = 2k^2 + k + 2$;
$B = k - 1$;

$C = 2k^2 - k + 2;$
$D = k + 1;$

$q = 3,\ m = 3k;$
$h = 27k^4 + 57k^2 + 27;$
$A = 3k^2 + k + 3;$
$B = k - 1;$
$C = 3k^2 - k + 3;$
$D = k + 1;$

Example 6.2.
$h = 8k^3p + 512k^7p^3 + 512k^3p^3 + 1024k^5p^3;$
$A = 8k^3p + 8kp - k;$
$B = k + 1;$
$C = 8k^3p + 8kp + k;$
$D = k - 1;$

$k = \frac{1}{2};$
$h = 100p^3 + p;$
$A = 10p - 1;$
$B = 3;$
$C = 10p + 1;$
$D = 1;$

Example 6.3.
$h = n(p^4 + (n^2 + 2)p^2 + 1);$
$A = p^2 + np + 1;$
$B = p + 1;$
$C = p^2 - np + 1;$
$D = p - 1;$

Remark 6. The case $n = 1$, recovers the last parametric solution of the table 1 in [A.C2].

Example 6.4. $q = 1;$
$h = m^4 + 3m^2 + 1;$
$A = m + m^2 + 1;$
$B = m - 1;$

$C = m - m^2 - 1;$
$D = m + 1;$

Again this recovers the last parametric solution of the table 1 in [A.C2].

Example 6.5.
$h = (p^2 + 2)(p^2 + 4);$
$A = p^2 + p + 2;$
$B = p + 1;$
$C = p^2 - p + 2;$
$D = p - 1;$

Example 6.6.
$h = 512m^4 + 1032m^2 + 512;$
$A = 8m^2 - m + 8;$
$B = m + 1;$
$C = 8m^2 + m + 8;$
$D = m - 1;$

Example 6.7. $q = m^2;$
$h = 1 + m^2 + m^6 + 2m^4;$
$A = 1 + m + m^3;$
$B = 1 - m;$
$C = 1 - m - m^3;$
$D = 1 + m;$

Example 6.8. By taking $Z = h^2 + 1$, in the elliptic curve (2.6) (or $X = 2h^2 + 1$ in the (2.5)), we get

$$Y^2 = h^6 - h^4 - h^2 + 1 = (h^2 - 1)(h^4 - 1) = (h^2 - 1)^2(h^2 + 1). \tag{6.6}$$

letting $h^2 + 1 = t^2$, yields $h = \frac{r^2-1}{2r}$ and $t = \frac{-(r^2+1)}{2r}$, then we get

$Y = (h^2 - 1)(-t) = \frac{(r^6-5r^4-5r^2+1)}{8r^3}$, and $X = \frac{r^4+1}{2r^2}$.

Finally by calculating m, p, q, A, B, C, D, from the above relations, we obtain the parametric solution as follows:
$h = 8r^3(r^2 - 1),$

$A = 4r(5r^4 - 1),$
$B = (r^4 + 6r^3 + 6r^2 + 6r + 1)(r - 1)^2,$
$C = 4r^3(r^4 - 5),$
$D = (r^4 - 6r^3 + 6r^2 - 6r + 1)(r + 1)^2.$

Note that h is given by polynomial of degree 5.

The Sage software and Denis Simon's ellrank code were used for calculating the rank of the elliptic curves, (see [S.A]).

Acknowledgements

We are very grateful to Professor Allen MacLeod for the rank and generator computations of the elliptic curves with big generators. Finally the second author would like to present this work to his parents and his wife.

References

[A.C] A. Choudhry, : On The Diophantine Equation $A^4 + hB^4 = C^4 + hD^4$, Indian J. Pure Appl. Math, $26(11), pp.1057 - 1061, (1995)$.

[A.C2] A. Choudhry, A Note on the Quartic Diophantine Equation $A^4+hB^4 = C^4+hD^4$, available at arXiv. (2016)

[L.D] L. E. Dickson, History of the Theory of Numbers, Vol. II: Diophantine Analysis, G. E. Stechert k Co., New York, (1934).

[T.P] T. Piezas, A collection of algebraic identities, available at https://sites.google.com/site/tpiezas/0021e, (accessed on 7 April (2016)).

[S.T] S. Tomita, https://www.maroon.dti.ne.jp/fermat/dioph121e.html, (accessed on 7 April (2016)).

[H.R] H. W. Richmond, On the Diophantine equation $a^4 + by^4 + cz^4 + dw^4 = 0$, the product abcd being a square number. J. Lond. Math. Soc., $19, 193 - 194.(1944)$.

[S.A] Sage software, available from http://sagemath.org.

[L.W] L. C. Washington, Elliptic Curves: Number Theory and Cryptography, Chapman-Hall, (2008)

Farzali Izadi, Department of Mathematics, Faculty of Science, Urmia University, Urmia 165-57153, Iran

E-mail address: f.izadi@urmia.ac.ir

Mehdi Baghalaghdam, Department of Mathematics, Faculty of Science, Azarbaijan Shahid Madani University, Tabriz 53751-71379, Iran

E-mail address: mehdi.baghalaghdam@yahoo.com